

A QUICK AND DIRTY IRREDUCIBILITY TEST FOR MULTIVARIATE POLYNOMIALS OVER $\mathbb{F}_q$

H.-C. GRAF V. BOTHMER AND F.-O. SCHREYER

ABSTRACT. We provide some statistics about an irreducibility/reducibility test for multivariate polynomials over finite fields based on counting points. The test works best for polynomials in a large number of variables and can also be applied to black box polynomials.

1. INTRODUCTION

Let $f \in \mathbb{F}_q[x_1, \dots, x_n]$ be a polynomial. Since $f(x)$ can take only q possible values for every point in $x \in \mathbb{A}^n(\mathbb{F}_q)$ we expect that $f(x) = 0$ for about $\frac{1}{q}$ of the points $\mathbb{A}^n(\mathbb{F}_q)$. If on the other hand $f = gh$ is a product of two polynomials $g, h \in \mathbb{F}_q[x_1, \dots, x_n]$, we have $f(x) = 0$ if $g(x) = 0$ or $h(x) = 0$. So one might expect that products of polynomials satisfy $f(x) = g(x)h(x) = 0$ for approximately $\frac{2}{q} - \frac{1}{q^2}$ of the points $x \in \mathbb{A}^n(\mathbb{F}_q)$. This phenomenon is well explained by the Weil formulas [Mil80].

In this article we investigate the following irreducibility test for multivariate polynomials f over $\mathbb{F}_q$:

Evaluate f at N random points. We reject the hypothesis that f is reducible, if the fraction of zeros $\gamma_q(f)$ found is significantly smaller than $\frac{2}{q} - \frac{1}{q^2}$. Note that 99.5% of all polynomial functions satisfy

$$\gamma_q(f) \leq \frac{1}{q} + 2.58 \sqrt{\frac{\frac{1}{q}(1 - \frac{1}{q})}{q^n}}.$$

This irreducibility test is quick, since the number of evaluations needed to detect a given percentage $1 - \epsilon$ of all products of polynomial functions or all general polynomial functions do not depend on the degree of the polynomials considered respectively, i.e.

$$N \sim O(-q \ln \epsilon).$$

On the other hand it is dirty, since it does not give a definite answer. Moreover we cannot make ϵ arbitrarily small, because N is bounded by q^n , the number of $\mathbb{F}_q$ rational points in $\mathbb{A}^n(\mathbb{F}_q)$. There will always be a few polynomials that cannot be correctly classified by our method at all. For example the product of an irreducible, absolutely reducible polynomial with a further absolutely irreducible polynomial.

Date: February 8, 2008.

The test works for implicitly given (black box) polynomials as well. We give examples of such polynomials below.

The expected fraction of zeros for special classes of polynomials is also larger than $\frac{1}{q}$. For example, the expected fraction of zeros for $n \times n$ determinants is

$$E(\gamma_{q,\det}) = 1/q + 1/q^2 - 1/q^5 - 1/q^7 + O(1/q^{12})$$

for $n \geq 12$.

Notation 1.1.

$\mathbb{F}_q$	the finite field with q elements
$X \subset \mathbb{A}^n$	an affine algebraic set
$X(\mathbb{F}_q)$	the $\mathbb{F}_q$ -rational points of X
$ X = X(\mathbb{F}_q) $	the number of $\mathbb{F}_q$ -rational points of X
$\gamma_q(X)$	the fraction of $\mathbb{F}_q$ -rational points in $\mathbb{A}^n$ that are contained in X
$\mathcal{B}(N, p, k) = \binom{N}{k} p^k (1-p)^{N-k}$	the binomial distribution
N	the number of trials
p	the success probability
k	the number of successes
$\mathcal{N}(\mu, \sigma)$	the normal distribution with mean μ and variance σ^2

$\mathcal{B}(N, p)$ can be approximated by $\mathcal{N}(p, \sqrt{p(1-p)/N})$.

2. FRACTIONS OF ZEROS

Example 2.1. We choose fixed polynomials f_1, f_2 of degree 5 and f_3 of degree 10 in $\mathbb{Z}[x_1, \dots, x_4]$ with coefficients in $[-9, 9]$ using the random number generator of the computer algebra system MACAULAY 2 [GS] and consider $f = f_1 f_2 + 7 f_3$. Let X be the vanishing set $V(f)$.

A black-box polynomial is a polynomial for which it is easy to check $f(x) = 0$, but the explicit formula for f in terms of the unknowns $x_1 \dots x_n$ is hard or impossible to write down.

Example 2.2. Let $S_d \subset H^0(\mathbb{P}^2, \mathcal{O}(d))$ be the hypersurface of singular homogeneous polynomials f of degree d in 3 variables. For each point $f \in H^0(\mathbb{P}^2, \mathcal{O}(d))$ it is easy to decide whether $f \in S_d$ via the Jacobi criterion. On the other hand the equation of S_d in the $\binom{d+2}{2}$ variables is not obvious.

Example 2.3. Let $C \subset \mathbb{P}^4$ be the determinantal curve of degree 10 and genus 6 defined by the maximal minors of the following 5×3 matrix

$$\begin{pmatrix} x_0 + x_1 - x_3 - x_4 & x_0 - x_1 - x_2 - x_4 & -x_0 + x_3 + x_4 \\ -x_0 - x_2 + x_3 + x_4 & x_0 - x_1 - x_2 - x_3 + x_4 & -x_0 + x_1 - x_2 + x_3 + x_4 \\ -x_0 - x_2 - x_3 - x_4 & -x_0 - x_1 - x_3 - x_4 & -x_1 + x_4 \\ -x_1 - x_2 - x_3 + x_4 & -x_1 - x_2 & -x_1 + x_2 \\ -x_0 + x_1 - x_2 - x_3 - x_4 & -x_0 + x_2 - x_3 + x_4 & x_0 - x_1 + x_2 + x_3 + x_4 \end{pmatrix}$$

Let $D = \{H \in \check{\mathbb{P}}^4 \mid H \cap C \text{ is singular}\}$ be the dual variety of C .

Definition 2.4. Let $X \subset \mathbb{A}^n$ an algebraic set. We denote by

$$\gamma_q(X) := \frac{|X(\mathbb{F}_q)|}{|\mathbb{A}^n(\mathbb{F}_q)|}$$

the fraction of $\mathbb{F}_q$ -rational points on X . In particular for a hypersurface $X = V(f)$ we have $\gamma_q(f) = \gamma_q(V(f))$. We call $\gamma_q(f)$ the fraction of $\mathbb{F}_q$ -rational zeros of f .

Example 2.5. We estimate γ_q in three of our examples by evaluating in $N = 1000$ random points over all primes up to 17. The following table gives the 99% confidence interval for γ_q :

q	X	S_8	D
2	$56.7\% \pm 4.0\%$	$68.4\% \pm 2.9\%$	$55.3\% \pm 4.1\%$
3	$33.8\% \pm 3.9\%$	$42.3\% \pm 3.1\%$	$49.2\% \pm 4.1\%$
5	$17.9\% \pm 3.1\%$	$24.0\% \pm 2.6\%$	$24.9\% \pm 3.5\%$
7	$26.2\% \pm 3.6\%$	$16.8\% \pm 2.3\%$	$35.3\% \pm 3.9\%$
11	$9.3\% \pm 2.4\%$	$8.9\% \pm 1.8\%$	$8.0\% \pm 2.2\%$
13	$8.6\% \pm 2.3\%$	$9.6\% \pm 1.8\%$	$8.4\% \pm 2.3\%$
17	$5.2\% \pm 1.8\%$	$8.1\% \pm 1.7\%$	$5.9\% \pm 1.9\%$

In this article we will explain these numbers.

Remark 2.6. We can compute the true values $\gamma_2(X) = 56.3\%$, $\gamma_3(X) = 34.6\%$, $\gamma_5(X) = 18.7\%$ and $\gamma_7(X) = 27.6\%$ with the same effort, since there are less than 1000 rational points in $\mathbb{A}^4(\mathbb{F}_q)$ for $q \leq 7$.

To study the map

$$\gamma_q: \mathbb{F}_q[x_1 \dots x_n] \rightarrow [0, 1], f \mapsto \gamma_q(f)$$

we note that $\gamma_q(f)$ factors over the ring $R := \text{map}(\mathbb{A}^n(\mathbb{F}_q), \mathbb{F}_q)$:

$$\begin{array}{ccc} \mathbb{F}_q[x_1 \dots x_n] & \xrightarrow{\gamma_q} & [0, 1] \\ \downarrow \psi & \nearrow & \\ R & & \end{array}$$

Lemma 2.7. ψ is surjective.

Proof. Since $|\mathbb{A}^n(\mathbb{F}_q)| = q^n < \infty$ we can find a polynomial with prescribed values at these points via interpolation. $\square$

We study the distribution of γ_q on R by regarding it as a random variable on the finite probability space

$$(R, \Omega, P)$$

with Ω the sigma algebra of all subsets of R and P the constant probability measure.

Proposition 2.8. The distribution of γ_q on R is binomial

$$P\left(\gamma_q = \frac{k}{q^n}\right) = \mathcal{B}\left(q^n, \frac{1}{q}, k\right).$$

In particular the expectation value of γ_q is $E(\gamma_q) = \frac{1}{q}$.

Proof. We have to count the maps $f \in R$ that map precisely k different points to 0. Since the values at different points are independent, this number is

$$\binom{q^n}{k} 1^k \cdot (q-1)^{q^n-k}$$

The probability that $\gamma_q = \frac{k}{q^n}$ is therefore

$$P\left(\gamma_q = \frac{k}{q^n}\right) = \binom{q^n}{k} \left(\frac{1}{q}\right)^k \cdot \left(\frac{q-1}{q}\right)^{q^n-k} = \mathcal{B}\left(q^n, \frac{1}{q}, k\right)$$

□

Example 2.9. Consider maps $f \in R = \text{map}(\mathbb{A}^4(\mathbb{F}_{11}), \mathbb{F}_{11})$. The distribution of fractions of zeros is

$$P(\gamma_{11} = k/11^4) = \mathcal{B}(11^4, 1/11, k).$$

From its approximation by the normal distribution $\mathcal{N}(0.0909, 0.0024)$ we obtain

$$P(0.0847 \leq \gamma_{11} \leq 0.0971) \leq 99\%.$$

We now consider products. The random variable

$$\gamma_{q,\cup}: R \times R \rightarrow [0, 1], \gamma_{q,\cup}(f, g) = \gamma_q(fg) = |V(f) \cup V(g)|/q^n$$

which assigns to each pair of functions the fraction of zeros of their product.

Proposition 2.10. *On $R \times R$ the distribution of $\gamma_{q,\cup}$ is*

$$P(\gamma_{q,\cup} = k/q^n) = \mathcal{B}(q^n, (2q-1)/q^2, k).$$

In particular the expectation value of $\gamma_{q,\cup}$ is

$$E(\gamma_{q,\cup}) = \frac{2q-1}{q^2} = 1 - \left(\frac{q-1}{q}\right)^2.$$

Proof. The value of $f \cdot g$ in a point x depends on the values of f and g at x . There are q^2 ways of choosing these values of which $(q-1)^2$ give $(f \cdot g)(x) \neq 0$. □

Example 2.11. Consider pairs (f, g) of functions in R as in Example 2.9. The distribution of $\gamma_{11,\cup}$ is now

$$P(\gamma_{11,\cup} = k/11^4) = \mathcal{B}(11^4, 21/11^2, k).$$

From its approximation by the normal distribution $\mathcal{N}(0.1736, 0.0031)$, we obtain

$$P(0.1655 \leq \gamma_{11,\cup} \leq 0.1816) \geq 99\%$$

Note that this range does not intersect

$$P(0.0847 \leq \gamma_{11} \leq 0.0971) \geq 99\%.$$

Geometrically products of functions correspond to the union of their zero-sets. γ_q also behaves well under other geometric operations:


Points on a hypersurface of degree 10 in $\mathbb{A}^4$ 

FIGURE 1. 99% of polynomial functions on $\mathbb{A}^4$ have γ_q between the continuous lines. 99% of products have γ_q between the dashed lines.

Proposition 2.12 (Intersection). *Let $X \subset \mathbb{A}^n$ be a subvariety. We consider the random variable*

$$\gamma_{q,\cap X}: R \rightarrow [0, 1], \quad \gamma_{q,\cap X}(f) = |V(f) \cap X|/q^n.$$

The distribution of $\gamma_{q,\cap X}$ is

$$P(\gamma_{q,\cap X} = k/q^n) = \mathcal{B}(|X|, 1/q, k).$$

In particular, the expectation value of $\gamma_{q,\cap X}$ is $E(\gamma_{q,\cap X}) = \gamma_q(X)/q$, where $\gamma_q(X) = |X|/q^n$ is the fraction of points on X in $\mathbb{A}^n(\mathbb{F}_q)$.

Proof. Clearly, $x \in X \cap V(f)$ if and only if $x \in X$ and $f(x) = 0$. Since the values of f can be chosen independently on the points of X , we have

$$P(x \in \ker f \cap X | x \in X) = \frac{1}{q}.$$

□

Corollary 2.13. *Consider the random variable*

$$\gamma_{q,\cap}: R^c \rightarrow [0, 1], \quad \gamma_{q,\cap}(f_1, \dots, f_c) = |V(f_1) \cap \dots \cap V(f_c)|/q^n$$

Then the expected fraction of points is $E(\gamma_{q,\cap}) = \frac{1}{q^c}$

Proof. Use Proposition 2.12 inductively. □

Notice that for polynomials $f_1, \dots, f_c$ the expected codimension of $V(f_1, \dots, f_c) \subset \mathbb{A}^n$ is also c .


Singular curves in $\mathbb{P}^2$ 

FIGURE 2. The diagram shows the expectation values for various classes of polynomials in a large number of variables, and the measurement for S_8 , the hypersurface of singular plane curves of degree 8. Note that the diagram tells that about 70 % of all plane curves over $\mathbb{F}_2$ are singular.

Proposition 2.14 (Substitution). *Let $R^m = \text{map}(\mathbb{A}^n(\mathbb{F}_q), \mathbb{A}^m(\mathbb{F}_q))$ and $X \subset \mathbb{A}^m(\mathbb{F}_q)$ a subset. Consider the random variable*

$$\gamma_{q, \text{subst}}: R^m \rightarrow [0, 1], \quad \gamma_{q, \text{subst}}(\phi) = |\phi^{-1}X|/q^n$$

The distribution of $\gamma_{q, \text{subst}}$ is

$$P(\gamma_{q, \text{subst}} = k/q^n) = \mathcal{B}(q^n, \gamma_q(X), k).$$

In particular the expectation value of $\gamma_{q, \text{subst}}$ is $E(\gamma_{q, \text{subst}}) = \gamma_q(X) = |X|/q^n$.

Proof. Choosing functions $f_1, \dots, f_n$ is equivalent to independent choice of the image points. Therefore the probability of $\phi^{-1}(X)$ containing exactly k points is the same as the probability of hitting k points of X while choosing q^n points in $\mathbb{F}_q^n$. This gives the desired binomial distribution. $\square$

3. DETERMINANTAL VARIETIES

Even though we have shown, that $E(\gamma_q) = \frac{1}{q}$ with a small variance on the set of all functions from A to $\mathbb{F}_q$, there are special classes of functions that have larger expected γ_q .

It turns out that this behavior is common for determinants:

Proposition 3.1. *Let $X \subset \mathbb{A}^{nm}$ be the determinantal variety of $n \times m$ matrices with $n \leq m$ of rank less than n . Then the fraction of points on X is*

$$\gamma_q(X) = 1 - \prod_{i=0}^{n-1} \left(1 - \frac{1}{q^{m-i}}\right),$$

i.e. X contains $\gamma_q(X) \cdot q^{nm}$ points.

Proof. We prove that the number of matrices that have maximal rank is

$$\prod_{i=0}^{n-1} (q^m - q^i)$$

by induction. M is a matrix of full rank if and only if the first $n-1$ rows form a matrix of full rank and the last row is linearly independent of the first $n-1$ rows. Since there are q^{n-1} linear combinations of the first $n-1$ rows we obtain a further factor $(q^m - q^{n-1})$.

□

Corollary 3.2. *On the space of matrices R^{nm} , consider the random variable*

$$\gamma_{q,\det}: R^{nm} \rightarrow [0, 1], \quad \gamma_{q,\det}(M) = |\{x \in \mathbb{A}^n \mid \text{rank } M(x) < n\}|/q^n.$$

Then the fraction of zeros has expectation value

$$E(\gamma_{q,\det}) = 1 - \prod_{i=0}^{n-1} \left(1 - \frac{1}{q^{m-i}}\right) = \frac{1}{q^{m-n+1}} + \dots$$

The distribution of $\gamma_{q,\det}$ is

$$P(\gamma_{q,\det} = k/q^n) = \mathcal{B}(q^n, E(\gamma_{q,\det}), k)$$

Proof. Substitute functions for the variables in the generic $n \times m$ matrix and use Proposition 2.14 □

In the special case of $n \times n$ square matrices we have

$$E(\gamma_{q,\det}) = 1/q + 1/q^2 - 1/q^5 - 1/q^7 + O(1/q^{12})$$

for $n \geq 12$.

Example 3.3 (Example 2.2 continued). For small primes the divisor S_d has more points than expected for irreducible polynomials, but not enough to seem reducible, see Figure 2. Our measurements are consistent with the well known fact that S_d is an irreducible determinantal hypersurface [GKZ94, Chapter 13, Prop. 1.6 and 1.7].

Points on the dual variety of a curve in $C \subset \mathbb{P}^4$




FIGURE 3. C has a simple node over $\mathbb{F}_7$ and is smooth over $\mathbb{F}_p$ for $p = 5, 11, 13, 17$.

4. TESTING

To decide between two binomial distributions with success probabilities $p_1 < p_2$ and N experiments, we compute empirical probability $\bar{p} = \frac{k}{N}$ and decide for p_1 if

$$\bar{p} \leq p_{middle} = \frac{\sqrt{p_1 p_2} (\sqrt{p_1(1-p_2)} + \sqrt{p_2(1-p_1)})}{\sqrt{p_1(1-p_1)} + \sqrt{p_2(1-p_2)}} \approx \sqrt{p_1 p_2}.$$

To achieve a confidence level of $1 - \epsilon$ we choose $s = s(\epsilon)$ such that

$$\Phi(s) = \frac{1}{\sqrt{2\pi}} \int_s^\infty e^{-\frac{x^2}{2}} dx = \epsilon$$

and N such that

$$\sqrt{N} \geq s(\epsilon) \frac{\sqrt{p_1(1-p_1)} + \sqrt{p_2(1-p_2)}}{p_2 - p_1}.$$

In our case we have

$$p_1 \leq \frac{1}{q} + s(\epsilon) \sqrt{\frac{\frac{1}{q}(1 - \frac{1}{q})}{q^n}}$$

for $1 - \epsilon$ of all polynomials and

$$p_2 \geq \frac{2q-1}{q^2} - s(\epsilon) \sqrt{\frac{\frac{2q-1}{q^2}(1 - \frac{2q-1}{q^2})}{q^n}}.$$

for $1 - \epsilon$ of all products of polynomials. The decision based on the empirical probability $\bar{p} = \frac{k}{N}$, is then correct in $1 - \epsilon$ cases of the experiments. Note

however, that for fixed n and q we cannot make ϵ arbitrarily small, since we need $p_1 \leq p_2$.

An easy calculation gives the following estimate

$$\sqrt{N} \geq s(\epsilon) \frac{(2q)^{\frac{3}{2}}}{q - 1 - 2sq^{-\frac{n-2}{2}}}$$

for $q \geq 3$, which approaches $s(\epsilon)\sqrt{2q}$ for large n or q . Since $s(\epsilon) = O(\sqrt{-\ln(\epsilon)})$, we conclude that N grows like $O(-q \ln \epsilon)$.

For $\epsilon = 0.5\%$, $s = 2.58$, the number of trials needed is

	2	3	5	7	11	13	17
$n = 1$	∞	∞	∞	∞	∞	∞	∞
$n = 2$	∞	∞	∞	∞	∞	∞	∞
$n = 3$	∞	∞	∞	28373	2355	1908	1669
$n = 4$	∞	∞	1103	647	634	682	803
$n = 5$	∞	1705	367	369	482	551	695
$n = 6$	∞	384	259	308	447	521	673
$n = 7$	4457	224	225	289	437	513	667
$n = 8$	619	173	212	283	434	511	666
$n = 9$	295	151	206	280	433	511	666
$n = 10$	197	140	204	279	433	511	665

∞ indicates that there are not enough points in $\mathbb{A}^n(\mathbb{F}_q)$ to perform the test for the required $\epsilon = 0.5\%$. In case we can perform the test, the deciding number of successes Np_{middle} is

	2	3	5	7	11	13	17
$n = 1$	∞	∞	∞	∞	∞	∞	∞
$n = 2$	∞	∞	∞	∞	∞	∞	∞
$n = 3$	∞	∞	∞	5607	301	207	139
$n = 4$	∞	∞	303	128	81	74	66
$n = 5$	∞	754	101	73	61	59	57
$n = 6$	∞	170	71	61	57	56	55
$n = 7$	2821	99	61	57	55	55	55
$n = 8$	391	76	58	56	55	55	55
$n = 9$	186	67	56	55	55	55	55
$n = 10$	125	62	56	55	55	55	55

5. HIGHER CODIMENSION

In principle this method can be applied to algebraic sets of higher codimension.

Consider two surfaces in $\mathbb{P}^4$ and their union. We would like to distinguish their union from the irreducible examples. One possibility is to consider the Chow form which is a determinantal hypersurface on $G(2, 5)$ in this case. In Figure 4 we indicate the 5% and the 95% quantiles of γ_q for the Chow forms of 100 Bordiga surfaces, elliptic scrolls and their unions. A second possibility is to count points and apply Corollary 3.2. As Figure 4 shows there is no difference between the two methods. The formula for the error


Surfaces in $\mathbb{P}^4$ 

FIGURE 4. The 5% and the 95% quantiles of γ_q for the Chow forms of 100 Bordiga surfaces, elliptic scrolls and their unions compared with the error estimates for counting points on codimension 2 determinantal varieties rescaled. Using the geometry of Bordiga surfaces we obtain a better estimate.

term underestimates the number of points on a elliptic scroll, because the scroll is irregular.

The method of searching points at random in higher codimensional subsets of rational varieties helped us in proving the existence of several interesting components of Hilbert schemes. [Sch96], [ST02], [vBEL04]

REFERENCES

- [GKZ94] I. M. Gel'fand, M. M. Kapranov, and A. V. Zelevinsky. *Discriminants, resultants, and multidimensional determinants*. Mathematics: Theory & Applications. Birkhäuser Boston Inc., Boston, MA, 1994.
- [GS] Daniel R. Grayson and Michael E. Stillman. Macaulay 2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2>.
- [Mil80] James S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980.
- [Sch96] Frank-Olaf Schreyer. Small fields in constructive algebraic geometry. In *Moduli of vector bundles (Sanda, 1994; Kyoto, 1994)*, volume 179 of *Lecture Notes in Pure and Appl. Math.*, pages 221–228. Dekker, New York, 1996.
- [ST02] Frank-Olaf Schreyer and Fabio Tonoli. Needles in a haystack: special varieties via small fields. In *Computations in algebraic geometry with Macaulay 2*, volume 8 of *Algorithms Comput. Math.*, pages 251–279. Springer, Berlin, 2002.
- [vBEL04] H.-Chr. Graf v. Bothmer, C. Erdenberger, and K. Ludwig. A new family of rational surfaces in $\mathbb{P}^4$. preprint, 2004.

INSTITUT FÜR MATHEMATIK (C), WELFENGARTEN 1, UNIVERSITÄT HANNOVER, D-30167
HANNOVER, GERMANY

E-mail address: `bothmer@math.uni-hannover.de`

MATHEMATIK UND INFORMATIK, GEB. 27, UNIVERSITÄT DES SAARLANDES, D-66123
SAARBRÜCKEN, GERMANY

E-mail address: `schreyer@math.uni-sb.de`